Recovery Kernel

DISCONTINUANCE PROTOCOL: METHAMPHETAMINE

Die Kras, die Tydlyn, en die Pad Terug na 'n Werkende Dopamienstelsel

---

This guide follows the same structure as the Recovery Kernel's Forced Discontinuance series — a safety framework, a genetic cross-reference, a use-pattern classification, a recovery protocol, and a full protocol map — adapted to what methamphetamine discontinuation actually is. It is not a seizure risk the way benzodiazepine withdrawal is. It is, in the Recovery Kernel's own words, 'one of the most profoundly unpleasant biological experiences' a person can go through, and it carries a real and serious risk of its own: the crash produces a depth of depression and anhedonia severe enough to be a genuine suicide risk window. This guide treats that risk with the same seriousness the benzo guide gives to seizures.

Section 0 — Immediate Safety: Read This First

⚠ WAARSKEUWING

METHAMPHETAMINEWITHDRAWAL WILL NOT KILL YOU THROUGH SEIZURE THE WAY BENZODIAZEPINE OR ALCOHOLWITHDRAWAL CAN. BUT THE CRASH THAT FOLLOWS HEAVY OR PROLONGED USE PRODUCES ONE OF THE MOST SEVERE PSYCHOLOGICAL STATES IN ADDICTION MEDICINE, AND THE DEPRESSION IT PRODUCES IS A GENUINE SUICIDE RISK WINDOW. TAKE THE CRASH AS SERIOUSLY AS ANY PHYSICAL WITHDRAWAL DESCRIBED ELSEWHERE IN THIS LIBRARY.

Seek immediate emergency help if you experience:

- As any thoughts of suicide or self-harm, or a sense that the depression is more than you can get through — call or text findahelpline.com, or text HOME to 741741, both available 24/7.
- Chest pain, irregular heartbeat, shortness of breath, sudden severe headache, or facial drooping/arm weakness — this can indicate a cardiac event or stroke, both real risks after heavy stimulant use, and require calling 112 immediately.
- Paranoid delusions, hallucinations, or an inability to distinguish what's real — meth-induced psychosis can persist for days to weeks after last use and requires medical evaluation.
- Ekstreme agitering of 'n gevoel van die verlies van beheer oor jou aksies

SANCTUARY

Die kras self – extreme uitputting, hypersomnie, intense honger en 'n golf van depressie – word verwag en is nie op sy eie 'n mediese noodgeval nie. Dit is die liggaam wat begin herstel. Die onderskeiding wat saak maak, is dieselfde as wat die SSRI-gids noem: gemoedstoestand en veiligheid, nie ongemak nie. Indien twyfel, gryp aan.

Section 1 — Hoekom Hierdie Gids Bestaan

The Recovery Kernel's own meth guide states this plainly: 'something has not gone wrong. Your brain is recalibrating. The misery is the signal that repair is beginning.' That sentence is the organising idea behind this entire discontinuance protocol. Meth withdrawal has no chemical antidote and no safe substitute-and-taper strategy the way benzodiazepines or opioids do — there is no equivalent of a cross-taper here. What there is instead is a well-documented, predictable biological timeline, and knowing that timeline in advance is itself a form of harm reduction: it is the difference between 'this misery means I need more meth to feeling normal' and 'this misery is Day 3 of a documented process that ends.'

'n Kort Geskiedenis: Hoe die Gevaar Ontdek Is

Methamphetamine is first synthesized from ephedrine in 1893 by the Japanese chemist Nagai Nagayoshi, and crystallised into the form still recognisable today in 1919 by Akira Ogata. It was patented and sold in Japan in the 1920s under the brand name Philopon, marketed for fatigue, narcolepsy, and weight loss — a legitimate pharmaceutical product, not yet understood as dangerous.

Dit het verander met sy massale uitrol tydens die Tweede Wêreldoorlog. Duitse chemici by Temmler het hul eie methamphetamine-formule in 1937 gepatenteer, en teen 1938 is dit onder die handelsnaam Pervitin oor die toonbank te koop — versprei aan Wehrmacht-soldate in enorme hoeveelhede (skattings bereik die tientalle miljoene tablette) spesifiek om die slapelose, geforseerde-mars tempo van die Blitzkrieg te onderhou, en aan siviele fabrieksarbeiders om lang skofte te onderhou. Japan het 'n parallelle program geërf, wat methamphetamine aan kamikaze-pilote en aan siviele fabrieksarbeiders vir dieselfde rede uitgereik: dit het uitgeputde mense laat funksioneer verby die punt their bodies would otherwise have stopped them.

DIE CENTURION

Dit is die historiese oorsprong van presies hierdie patroon in Afdeling 5 van hierdie gids — om 'n stimulerend middel te gebruik om 'n onhoudbare werktempo te onderhou. Dit was waar vir uitgeputte soldate en fabriekswerkers in 1940, en dit is vandag ook waar vir dieselfde patroon.

The danger became undeniable in the aftermath. Japan in particular experienced a massive postwar epidemic once wartime military stockpiles were dumped onto the civilian market — by some estimates, a meaningful percentage of the population was misusing the drug within a few years of the war's end. The scale of addiction, psychosis, and physical breakdown this produced was severe enough that Japan enacted the Stimulant Control Law in 1951, one of the first drug-specific stimulant regulations in the world — a direct legislative response to what unrestricted access had just demonstrated at national scale. What took a global war and 'n nasionale epidemie om te vestig in die 1940's is, in miniatuur, dieselfde les wat hierdie gids se Afdeling 2 tydlyn bestaan om vooraf te leer: hierdie dwelm se vermoë vir skade skaal direk met hoeveel die liggaam gevra word om sonder rus op dit te loop.

Section 2 — Die onttrekkingslyn

Timeline	What's Happening Biologically	Wat dit voel soos	What to Do
Hours 0–24 (Die Kraak)	Dopamine reserves exhausted; receptor signalling collapsing	Extreme moegheid, oormatosie, intense honger, depressie, angs	Slaap. Eet. Hidreer. Moenie die kras teenstryd nie – dit is die liggaam wat begin om te herstel. Elektrolyte.
Days 1–5 (Die Put)	Dopamine op die vloer; D2 reseptore nog ondergereguleer; die stelsel skree effektief na die chemikalie waaraan dit aangepas het.	Swere depressie, anhedonie (onvermoë om plesier te voel), diepgaande vermoeidheid, intense hunkeringe, versteurde slaap, kognitiewe mistigheid	Dit is die hoogste terugval en hoogste suïsidestrisiko venster in hierdie hele gids. Moenie alleen wees indien vermybaar nie. Stuur na findahelpline.com indien krisis. Begin met die Tier 1 aanvullingsondersteuning (Afdeling 8).
Days 5–14 (Die Grijs)	Acute symptoms resolving; dopamine system still significantly impaired; receptor recovery barely beginning	Constant flatness, low motivation, ongoing insomnia, cognitive impairment, intermittent cravings	Start with some light exercise—this is the repair signal (Section 3). Begin identifying and avoiding environmental use-triggers.

Timeline	What's Happening Biologically	Wat dit voel soos	What to Do
Weeks 2–4 (Die Stadige Terugkeer)	DRD2 receptor density begins measurable recovery; dopamine synthesis slowly increasing; prefrontal cortex repair initiating	Gradual, uneven mood improvement; cognitive fog lifting; cravings becoming episodic rather than constant; sleep stabilising	Establish a consistent daily routine. The first meaningful behavioural goals become realistic here, not before.
Maande 1–3	Measurable DRD2 receptor recovery; grey matter rebuilding in the prefrontal cortex; dopamine synthesis normalising	Genuine improvement with periodic cravings and mood variability; post-acute withdrawal (PAWS) episodes possible	Identity reconstruction work becomes appropriate here — not earlier. This is when capacity genuinely starts returning.
Maande 3–12	Significante reseptorherstel; BDNF-gedreven neuroplasticiteit herbouwt aktief beskadigde dopaminaxonen; functie van die prefrontale korteks aansienlik herstel.	Mood near baseline for most people; cravings are mostly context/cue-triggered rather than spontaneous; cognitive function largely restored; genuine enjoyment of life becomes accessible	Maintain every protocol below. Protect this window from major stress events where possible — the system is rebuilding, not yet at full structural strength.

Section 3 — Hoe lank voordat die brein regtig normaal is

This is die vraag onder elke fase van die tydlyn hierbo, en dit verdien 'n direkte antwoord eerder as 'n vae gerusstelling. Die eerlike antwoord het twee dele: die akute krisis los binne weke op, maar volledige reseptor-vlak herstel neem aansienlik langer — en om die werklike getal te ken verander wat "normaal" behoort te voel by elke fase.

DIE CENTURION

Die spesifieke datapunt, gesit direk uit die neurobeeldingsonderzoek waarop die Recovery Kernel se eie materiaal gebaseer is (Volkow et al.): dopamien D2 reseptorherstel na methamfetamiene-stop is beduidend, maar substansiële herstel neem 12 tot 17 maande volgehoue abstinentie. Dit is meetbaar stadiger as alkohol (beduidende D2 reseptorverbetering teen maand 1, volgens Heinz et al.) of kokain — methamfetamiene veroorsaak die mees uitgebreide dopamineterminaaldegawe van die algemene stimulerende middele, en die hersteltydlyn weerspieël dit.

Binne daardie groter 12-tot-17 maande boog, is daar 'n vinniger, meer akute herbouvenster wat spesifiek bekend gemaak moet word: die eerste 90 dae. Die

Recovery Kernel se neuroplastisiteitsraamwerk dokumenteer dit in detail — dae 31 tot 60 is wanneer nuwe neuraal paaie begin vorm en oues swak word; dae 61 tot 90 is wanneer daardie nuwe patrone begin konsolideer na struktureel meetbare verandering. Teen 90 dae, met konsekwente oefening, slaap, en die ander neuroplastisiteitsinvoer in Afdeling 8 hieronder, word dopamine reseptordigtheid aansienlik verbeter, grijsmaterievolume is meetbaar verhoog, en hunkeringe word dramatisch verminder vir die meeste mense — even though the full 12-17 month recovery is still in progress underneath that.

Recovery Milestone	Realistic Timeline
Acute crash resolves	Days 1–5
Cravings shift from constant to episodic/cue-triggered	Weeks 2–4
First measurable DRD2 receptor recovery on imaging	Weeks 2–4 through Month 1
Die 90-dae neuroplastisiteitsvenster sluit – nuwe gewoonte-argitektuur struktureel gekonsolideer.	Day 90
Near-baseline mood and cognition for most people	Maande 3–12
Substantial DRD2 receptor recovery (die hoofgetal)	12–17 months

BEING HEARD

As jy in maand 4 is en steeds plat, griewige dae het, beteken dit nie dat die proses misluk het of dat jy op enige manier gebreek is wat die tydlyn nie rekening hou mee nie. Volgens die data hierbo, is jy binne 'n werklik nog-herstel stelsel — 'naby-basislyn' by maande 3-12 beteken presies dit: nader, nie klaar nie. Die eindstreep vir die diepste laag van hierdie herstel is nader aan 'n jaar en 'n half as teenoor 'n seisoen. Dit is nie 'n ontmoedigende nommer nie. Dit is 'n eerlike een, en dit beteken dat die plat dae wat jy nog het, verwag word, nie bewys van mislukking nie.

'n Eerlike Waarskuwing: Hierdie tydlyne gaan uit van ware afstinensie.

Elke getal in hierdie afdeling – die weke, die maande, die 12-17 maand hoofgetal – is gemeet in mense wat met methamphetamine gestop het en daar gebly het. In die praktyk lyk baie herstel nie so skoon nie. Twee vervanginge is algemeen genoeg om spesifiek te noem, want elkeen interaksieer met presies die herstelmeganismes wat hierdie afdeling beskryf, en elkeen vertraag waarskynlik die klok eerder as om 'n neutrale brug dauroo te wees.

Swap out	What It Actually Does to the Recovery Mechanism	Likely Effect on the Timeline
Caffeine (used to manage the fatigue and low-motivation phases in Section 2)	Caffeine doesn't deplete dopamine the way methamphetamine does, so it isn't 'the same thing in miniature.' But it does stimulate the HPA axis — adding cortisol-system activation on	It plausibly prolongs the timeline indirectly, by undermining the sleep quality the actual repair work depends on, rather

Swap out	What It Actually Does to the Recovery Mechanism	Likely Effect on the Timeline
	top of a system, per Section 6, that may already be running hot (especially in the Met/Met 'Worrier' profile) — and heavy or late-day caffeine use directly interferes with sleep quality and duration. Section 8's Tier 3 is explicit that sleep is when BDNF synthesis and dopamine-system repair peak.	than by any direct dopamine mechanism. The more caffeine is substituted specifically to power through the fatigue Section 2 describes, the more it's fighting the exact process that fatigue is part of.
Alcohol (used to fall asleep)	This is the more concerning substitution of the two. Alcohol is sedating, but it fragments sleep architecture and specifically suppresses REM sleep — and REM is, per this library's own material, exactly the sleep stage where 'emotional memory processing and dopamine system maintenance occur.' Using alcohol to force sleep onset while suppressing the sleep stage recovery most depends on is close to solving the visible problem (can't fall asleep) while worsening the invisible one (the sleep that was supposed to repair the dopamine system isn't actually happening).	Plausibly prolongs the timeline directly, through degraded REM sleep, and introduces a second risk this section's numbers don't account for at all: a new dependence forming on top of the one being recovered from — genuinely likely for the DRD2 A1 allele / Reward Deficiency Syndrome profile in Section 6, where the reward circuit is already primed to over-attach to whatever substance is reliably available.

⚠ WAARSKEUWING

Neither of these substitutions is framed here as a moral failure — reaching for caffeine or a drink to get through an unbearable stretch is an extremely understandable response to real suffering. But the honest version of this guide has to say plainly: neither one is a neutral bridge through the timeline in the table above. If sleep is the problem, the non-alcohol tools in Section 8's Tier 3 (melatonin, magnesium glycinate, glycine, consistent schedule) are working with the repair mechanism instead of against it.

Die Breuk: Hoekom 'E Jaar Af, Dan Sal Ek Reg Wees' Nie Heeltemal Korrek Is Nie

There's an idea worth addressing directly, because it's a reasonable-sounding conclusion to draw from Section 3's own numbers, and it's not entirely wrong — which is exactly what makes it dangerous. The idea: 'if D2 receptors substantially recover by 12-17 months, I could take a year off, let my brain heal, and then know how to use safely again, or resume with a clean slate.' The old addiction-treatment

framing — 'the moment you pick up again, you're instantly back to full addiction' — isn't 100% true. But the instinct that a long break resets you to zero isn't 100% true Either. Die eerlike antwoord sit in die middel, en 'n frakatuur is 'n regtig nuttige manier om te beeld waarom.

What's True (Like a Healed Fracture)	What Else Is True (Hoekom Het Die Gang Nog Verander)
Die been genees. D2-reseptor-digtheid, volgens die data in Afdeling 3, herstel werklik op 'n regte, gemesuleerde tydlyn — dit is nie 'n eufemisme of 'n hoopvolle storie nie, dit is wat die beelde wys.	'n Geneesde been is nie dieselfde as die ongebreekte een nie. Dit herstruktureer rondom die besering. Dit kan weer werklike gewig dra — maar nie outomaties dieselfde las, op dieselfde hoek, op dieselfde manier, sonder dat die liggaam moet leer hoe nie.
Tolerance is genuinely lost during real abstinence — this is well-documented pharmacology, not guesswork.	Lost tolerance means a system that used to handle a given dose can no longer handle that same dose safely. Returning to a 'known' amount after a real break is dosing a body whose actual capacity has changed, even though the memory of what used to feel normal hasn't. This is precisely the mechanism behind the elevated overdose and cardiac-event risk (Section 0) seen specifically in relapse after a period of abstinence — not because the drug got stronger, but because the body's capacity to handle it got smaller while the expectation didn't.
Die ou "onmiddellike terugval na volle verslawing"-raamwerk oordry hoe vas en permanent die neurologie is.	It's not baseless, either. Section 6's epigenetic material describes how reward-circuitry gene expression shifts with repeated use — and those shifted pathways don't necessarily need to be rebuilt from nothing to reactivate. A previously-carved path can be walked again faster than a new one gets carved the first time. This is why relapse after a break often escalates in severity faster than the original use did, even when baseline receptor measures have genuinely recovered.

⚠ WAARSKEUWING

To be plain: a period of real abstinence is not a reset button, and it's not useless either — both of the framings the introduction to this section named are partly right and partly wrong at the same time. What it actually produces is a system that has repaired in the ways Section 3 describes, while remaining structurally different from the one that existed before use ever started — different tolerance, different vulnerability to fast reactivation, different cardiovascular capacity. Trying to put the exact same weight on it, the exact same way, is where the danger concentrates. This isn't a technical Note. It's the reason relapse after a real break carries its own distinct, elevated risk profile rather than being either 'safe now' or 'instantly back to square one.'

A Popular Claim Worth Correcting: 'Your Cells Fully Replace Every 7 Years'

This idea circulates widely, including inside recovery communities specifically — but it doesn't hold up to what's actually been measured, and the corrected version is more useful than the myth, not less. Different tissues turn over at wildly different rates: skin and gut lining in days to weeks, fat cells at roughly 10% per year, the skeleton over roughly a decade. But the tissue that matters most for everything in this guide — neurons in the cerebral cortex — are not meaningfully replaced at all after early development. The neurons involved in the reward pathways this entire guide describes are, To a large extent, they're the same physical cells you had before you ever used methamphetamine. They aren't swapped out for fresh ones on any timeline, seven years or otherwise.

What about age and time then? Two real things, distinct from the myth. First, what recovers on the timelines in this guide is not new cells replacing old ones — it's the density, sensitivity, and expression of receptors on the neurons that were already there, plus the synaptic connections between them (Section 6's epigenetics material). Second, aging does add a genuine complicating layer, through a different and well-documented mechanism: cellular senescence. As the body ages, some cells — including supporting glial cells in the brain — stop dividing but remain metabolically active, secreting inflammatory signalling molecules that accumulate over time and contribute to background neuroinflammation. This is real, current addiction and aging science, and it means the recovering system isn't a fixed target — it's ageing in the background at the same time it's repairing, which can plausibly make the whole process feel harder or slower the more time and the more cumulative use has passed, independent of the dopamine-specific timeline in Section 3.

BEING HEARD

because die dieselfde neurone bly bestaan eerder as dat hulle vervang word, die ou paaie wat hierdie gids bespreek het — die vinnige reaktivering risiko, die geleerde aanwysingsreaksies — is nie weg na 'n jaar af nie. Hulle is steeds fisies teenwoordig, in dieselfde selle, net stiller. Dit is presies hoekom om 'n jaar van regte werk af te skryf as 'dit het nie gewerk nie' of 'ek was nie sterk genoeg nie' of om na 'een keer 'n verslawende, altyd 'n verslawende' te gryp die verkeerde gevolgtrekking is wat gemaak kan word uit 'n terugval. — but so is assuming a year was ever going to erase pathways that were never going to be deleted in the first place, only outcompeted. The actual task was never full erasure. It's building new, competing pathways strong enough that the old ones stop being the path of least resistance — which is precisely why Section 11's Protocol 20/C-28 material talks about redirecting this exact reward circuitry toward genuine, high-intensity natural alternatives, rather than asking the circuitry itself to disappear. And it's worth saying plainly: returning to a substance after time away is not always someone deciding they could 'get away with it.' Gewone menslike nuuskierigheid – wonder, toets, bloot aangetrek tot iets op die manier waarop

mense na dinge aangetrek word – is 'n ware en verstaanbare trek op sy eie, sonder om 'n storie oor wilskrag of morele mislukking eraan te koppel.

Section 4 — Wat jy waarskynlik sal ervaar

Die tydlyn in Afdeling 2 vertel jou wat op biologiese vlak gebeur en omtrent wanneer. Hierdie afdeling is anders — dit is 'n poging om werklik te beskryf hoe dit voel van binne, meer diepgaand as wat 'n tabelry kan hou, sodat jy herken wanneer dit gebeur eerder as om paniekerig oor dit te wees.

Anhedonia — die platste woord vir die moeilikste simptoom

Anhedonia gets mentioned almost clinically in most withdrawal material — 'inability to feel pleasure' — as if that fully captures it. It doesn't. What it actually feels like, for most people, is closer to watching your own life through glass: food doesn't taste like much, music you used to love registers as sound without the feeling attached, a joke lands as information rather than something funny, a person you love doesn't produce the warmth being near them used to produce. It is not that you've stopped caring. It's that the machinery that used to translate 'good thing happening' into 'felt Good' is running at a fraction of its normal signal, because that machinery is the exact dopamine system methamphetamine hijacked and depleted. This is genuinely one of the most disorienting parts of the entire process, because it can produce a second-order fear on top of the flatness itself — 'what if I never feel anything again' — and that fear is worth naming directly: per Section 3's data, this is a documented, temporary state with a known recovery arc, not a permanent personality change.

Other symptoms worth naming plainly

- Hypervigilance and a racing, unsettled quality even through the exhaustion — the nervous system doesn't downshift as fast as the body needs sleep.
- Intense, specific cravings that come on with hardly any warning, often set off by something totally ordinary—a smell, a bit of tunes, a text ping—and that can feel, right then and there, like the only thing that'll make the present moment bearable.
- Irritability and a short fuse that can surprise you and the people around you, especially in the Pit phase (Days 1-5).
- Actual cognitive fog — word-finding difficulty, trouble following chats, a feeling like you're thinking through muck. This isn't a sign of permanent damage; it follows the same recovery curve as everything else in Section 2 and 3.
- Vivid, sometimes disturbing dreams once sleep starts returning — the brain doing a large amount of catch-up REM processing after a period of suppressed or fragmented sleep.

This one is worth a closer look, because it's often the most confusing symptom rather than the most obviously distressing one. The same BDNF-driven synaptic strengthening that's repairing the dopamine system (Section 3) works partly through glutamate signalling — glutamate and NMDA receptor activity are a normal, necessary part of how synapses actually get stronger, not purely a toxic byproduct. So the intensified dream activity in this recovery window is best understood as dual causation: real repair work happening, with unusually vivid and frequent dreaming as a side effect of that work, not primarily a sign of ongoing damage. For some people this goes further than just vivid — dreaming intensely enough to physically wake up from it, sometimes repeatedly through the night, which becomes its own sleep-disruption problem layered on top of the insomnia already present. Others notice they're suddenly recalling dreams in detail they normally wouldn't — waking up and being able to describe a dream at length, sometimes catching themselves talking about it as though it were something that actually happened, before the distinction settles back in. Both of these are consistent with the REM rebound and synaptic remodeling happening during this window, not signs that something has gone wrong with your grip on reality.

BEING HEARD

As jy die beskrywing van anhedonie hierbo lees en dink 'dis presies dit, en ek het nie 'n woord vir dit gehad nie' — daardie reaksie is algemeen, en dit is die moeite werd om daarop te stop. Baie mense gaan deur hierdie spesifieke simptoom gevoel soosof iets uniek verkeerd met hulle is, veral omdat niemand dit genoeg gedetailleerd beskryf het vir hulle om dit te herken as 'n bekende, gedokumenteerde ding wat gebeur aan byna almal wat deur dit gaan nie. Dit het 'n naam. Dit het 'n tydlyn. Dis nie jou nie.

Wat Maak Dit Moeilik: Die Dinge Wat Bewustelik Vermy Word

Some of what happens during this recovery is going to happen no matter what — the timeline in Section 2 and Section 3 is largely fixed biology. But some of what makes the experience feel unbearable enough to drive someone back to using isn't the underlying repair process itself — it's specific, avoidable things stacked on top of it that make an already-hard window meaningfully harder. Knowing which is which matters, because it's easy to mistake 'this got worse because of what I ate/drank/didn't sleep' for 'this recovery isn't working,' when the actual answer is often much more fixable.

What Makes It Worse	Hoekom
Hoësuiker en hoëglisemiese kos	Bloedsuikerspieke en -krasse vererger die stemmingwisseling en energie-krasse wat reeds plaasvind as gevolg van dopaminedepletasie – 'n kras bo-op 'n kras.
Caffeine en alkohol	Fully covered in Section 3 above — caffeine indirectly, through sleep interference and HPA-axis stimulation; alcohol directly, through REM suppression on top of the REM rebound already happening.

What Makes It Worse	Hoekom
Nicotine and other stimulants	Die duidelike een, maar dit is die moeite om net te sê: enige stimulan, insluitend nikotiin, hou dieselfde beloning- en opwindingkringloop ingespan wat hierdie hele herstel probeer laat vestig.
Slechte slaapgewoontes – skerms laat in die nag, onstabiele slaaptye.	Dit ondermyn direk die slaapafhanklike herstelmeganisme (BDNF-syntese, geheugenkonsolidasie) wat Afdeling 8 se Vlak 3 identifiseer as die primêre drywer van die hersteltydlyn in Afdeling 3.
Dehydrasie en gemiste maaltye	Die brein wat hierdie baie herstelwerk moet doen, benodig konstante glukose en hidrasie; op leeg loop addere 'n tweede, onnodige laag van uitputting bo-op die biologiese een wat reeds plaasvind.
Langdurige isolasie	Nie 'n kos of substansie nie, maar dit is die moeite werd om hier by te sluit — isolasie vererger meetbaar elke onttrekkingssindroom wat hierdie biblioteek dokumenteer, en dit neem die ondersteuningskontak weg wat Afdeling 0 identifiseer as beskermend gedurende die hoogste-risiko venster.

SANCTUARY

Exercise is die duidelike uitsondering op hierdie lys, en verdien om genoem te word as een eerder as om saamgevat te word met waarskuwings oor te veel doen. Volgens Afdeling 8 se Tier 3, is dit die primêre meganisme wat BDNF genereer wat die hele herstellyn in Afdeling 3 dryf — meer daarvan, nie minder nie, is die algemene rigting om na te beweeg sover kapasiteit toelaat.

BEING HEARD

Here's the part worth saying directly: almost everything painful in this recovery — the flatness, the fatigue, the intense dreams, the cravings — is your body actively repairing itself. It is not a sign that something is wrong. It's easy, in the middle of it, to mistake the discomfort of repair for evidence that recovery isn't working, and to reach for the drug specifically to make an idiopathic-feeling wave of symptoms stop — not because the craving itself got stronger, but because nobody told you that what you were feeling was expected, temporary, and productive. Knowing that in advance doesn't Make the discomfort disappear. But it changes what the discomfort means, and that changes what you do about it.

Section 5 — Die Werkgedrewe Gebruikspatroon en die Droom-Waak Oorlapping

Nie almal wat metamfetamine-afhanklikheid ontwikkel het, het begin by 'n partytjie of 'n krisis. 'n Regte en algemene patroon is gebruik wat begin het, of vererger het, spesifiek om te werk — lang skofte, meervoudige werksplekke, sorgdrawe bo-op 'n volle werksdag, die soort chroniese oorbelasting waar ophou

nie voel soos 'n opsie nie. Metamfetamine se stimulerende effek kan vir 'n rukkie lyk as die enigste manier om aan 'n rooster te hou wat van die begin af nooit volhoubaar was nie. Hierdie patroon verdien sy eie vermelding omdat dit 'n spesifieke en aparte simptoom veroorsaak wat ander beskrywings van meth-gebruik dikwels heeltemal mis.

⚠ WAARSKEUWING

Chronic, severe sleep deprivation — the kind that develops from using a stimulant to push through the sleep the body was supposed to get — can produce a phenomenon where dream content begins intruding into waking perception: brief, vivid, dreamlike images or sensations while you are ostensibly awake, a sense of unreality, or moments where you can't quite tell whether something happened or you dreamed it. This is a well-documented consequence of severe sleep deprivation itself (sometimes called hypnagogic intrusion or microsleep with dream content), not a sign of psychosis on its own — though it shares some features with die psigose-risiko genoem in Afdeling 0, en enige weergawe hiervan wat paranoïese wanpersepsies of 'n breuk van werklikheidsbeoordeling insluit (in plaas van kort, droomagtige perseptuele intrusies) moet op dieselfde manier geëvalueer word as wat Afdeling 0 beskryf.

Asseblief, as hierdie patroon op jou van toepassing is, sluit die herstelopdrag iets in wat die res van hierdie gids nie ten volle dek nie: eintlik die onderliggende oorwerk oplos, en nie net die substansie nie. 'n Dopamine-stelsel herbou (Afdeling 3) terwyl jy terugkeer na dieselfde onhoudbare rooster wat daartoe bygedra het om die afhanklikheid in die eerste plek te skep, is herbou in dieselfde toestande wat dit gebreek het. Dit is werd om eerlik te noem eerder as om dit geïmpliseer te laat.

Section 6 — Genetiese Kruisverwysing

Gene	Wat dit regeer	Relevant here
COMT (Val158Met)	Dopamine clearance speed	Val/Val ('Warrior,' fast clearance) is die primêre kwesbaarheidsprofiel vir metamfetamiën spesifiek — die vinnig-klere, intensiteit-soekende stelsel wat in meth se ekstreme dopamiensein te vrede gebring het by die eerste ervaring wat dit ooit ten volle bevredig het. Herstel vereis ware hoë-intensiteit alternatiewe, nie net afstinensie nie.
DRD2 / ANKK1 (Taq1A, A1 allele)	DRD2 receptor density	Die A1-allel drager se beloningskringe loop met verminderde baselinie D2-digtheid – Beloningtekortsyndroom. Dit is die enigste mees belangrike genetiese teiken vir herstelsupplementasie (die 'Brain Stack' in Afdeling 8), aangesien dit die

Gene	Wat dit regeer	Relevant here
		reseptorstelsel self is wat sy digtheid moet herbou, en nie net die dwelm skoonmaak nie.
MAOA (uVNTR)	Dopamine, serotonin, and norepinephrine are neurotransmitters that play crucial roles in regulating mood, arousal, attention, and motivation. While they all function as chemical messengers across the synapses, their breakdown mechanisms and associated enzymes differ somewhat. **General Breakdown Pathways:** The primary way these neurotransmitters are cleared from the synaptic cleft is through reuptake by the presynaptic neuron or glial cells, followed by enzymatic degradation within the cell. * **Dopamine (DA):** * **Primary Enzyme:** Monoamine Oxidase (MAO) and Catechol-O-Methyltransferase (COMT). * **Mechanism:** Dopamine is a catecholamine, making it susceptible to breakdown by enzymes that target aromatic rings. MAO oxidizes dopamine, while COMT methylates the molecule, which facilitates its subsequent degradation. * **Speed Factors:** The activity and expression levels of MAO (especially MAO-A) and COMT are major determinants of how quickly DA is cleared from the synapse.	Lae-aktiewe allele laat monoamine akkumuleer; hoë-aktiewe allele put dit vinnig uit. Die tweede patroon word geassosieer met groter impulsiwiteit en gebruikernernigheid — relevant vir beide aanvullingstyd en hervalrisiko-bewustheid.

Gene	Wat dit regeer	Relevant here
	* **Serotonin (5-HT):** * **Primary Enzyme:** Monoamine Oxidase (MAO). * **Mechanism:** Serotonin is primarily broken down by MAO, particularly in the gut and central nervous system. It can also be metabolized via different pathways depending on its precursor availability. * **Speed Factors:** The activity of MAO dictates the rate of serotonin clearance. * **Norepinephrine (NE):** * **Primary Enzymes:** Monoamine Oxidase (MAO) and Catechol-O-Methyltransferase (COMT). * **Mechanism:** Like dopamine, NE is a catecholamine and is broken down by both MAO and COMT. * **Speed Factors:** The combined action of these two enzymes determines the clearance rate. **Summary Comparison:** | Neurotransmitter | Primary Degrading Enzymes | Key Metabolic Pathway | Relative Speed Determinants | | :--- | :--- | :--- | :--- | | Dopamine (DA) | MAO, COMT | Methylation and Oxidation | Activity of both MAO and COMT. | | Serotonin (5-HT) | MAO | Oxidation | Primarily dependent on MAO activity. | | Norepinephrine (NE) | MAO, COMT | Methylation	

Gene	Wat dit regeer	Relevant here
	and Oxidation | Combined action of MAO and COMT. | **Clinical Relevance:** The understanding of these breakdown pathways is critical because many antidepressant and antipsychotic medications work by inhibiting the enzymes responsible for clearing these neurotransmitters (e.g., SSRIs inhibit serotonin reuptake/breakdown; MAO inhibitors block the enzyme itself).	
BDNF (Val66Met)	Neural repair and plasticity signalling	Met carriers secrete less BDNF in response to activity. Exercise-driven BDNF increase is the primary neuroplasticity tool in this entire guide — it works regardless of genotype, but Met carriers may need higher exercise intensity to generate an equivalent repair signal. This is not optional in this recovery — it's the repair mechanism itself.
CYP2D6 / CYP2B6	Metabolism of certain recovery-relevant medications (e.g., bupropion)	Relevant if any medication is introduced during recovery — poor metabolisers risk toxicity at standard doses; ultra-rapid metabolisers may get no effect. Worth disclosing to a prescriber before starting anything.

Hoe COMT Status Werklik Verander Wat Die Crash Voel So

Die Dopamine-Transmitter en die Rol van COMT

COMT (Catechol-O-Methyltransferase) is 'n ensiem wat help om dopamien te metaboliseer. Dit speel 'n sleutelrol in hoe ons brein dopamien, 'n belangrike neurotransmitter, hanteer. Dopamien is betrokke by baie funksies, insluitend beweging, gemoedstoestand en beloning.

Wat gebeur wanneer COMT oormatig aktief of onvoldoende werk?

Die verhouding tussen COMT-genvariasie en dopamienvlakke:

**** **Hoë COMT-aktiwiteit:** As die ensiem baie goed werk, kan dit dopamien vinniger afbreek. Dit beteken dat jou brein se dopamiinniveaus vinnig kan daal na 'n laer vlak.***
**** **Lae COMT-aktiwiteit:** As die ensiem minder aktief is, kan dopamien langer in die systeem bly, wat soms tot oormatige of onstabiele niveaus kan lei.***

Hoe dit die "crash" beïnvloed:

Die gevoel van 'n "crash" word dikwels geassosieer met 'n skielike en merkbare val in energie, gemoedstoestand of fokus – dikwels na gebruik van stimulerende stowwe of na intense opwinding.

1. **Vinniger Afbraak (Hoë COMT):** As jou brein dopamien vinnig afbreek, kan die oorgang van 'n hoë vlak na 'n lae vlak baie skerp en dramatisch wees. Dit kan voel soos 'n harde val of uitputting wat vinniger kom as wat jy verwag het.
2. **Langtrangereffek (Lae COMT):** Hoewel dit nie altyd die "crash" veroorsaak nie, kan onstabiele dopamiinniveaus lei tot 'n gevoel van gemoedstoestand wat makliker te manipuleer is of waar die afname meer geleidelik maar aanhoudend voel.

Die Implikasie vir Bestuur:

Dit beteken dat mense met verskillende COMT-profiele potensieel verskillende reaksies op chemiese en emosionele stimuli kan hê. Die begrip van hierdie biologiese variasie help om te verstaan hoekom sommige mense 'n baie skerp "crash" ervaar, terwyl ander dit meer

geleidelik voel.

Belangrike Notas:

**** Hierdie is algemene inligting en vervang nie professionele mediese advies nie.***
**** As jy probleme met jou gemoedstoestand of energie vlakke opmerk, praat asb. met 'n gesaghebbende.***

Die COMT-ry tevore noem die genotiipes; dit gaan daaroor hoe dit eintlik dag tot dag lyk, want Val/Val en Met/Met trek so baie anders terug dat dit die moeite werd is om dit afsonderlik te beskryf eerder as om dit as 'n abstrakte label te laat.

COMT Profile	Baseline Before Meth	How the Crash Manifests
Val/Val ('Warrior') — vinnige dopamienklaring, laer basisdopamiennive au	Dikwilig om te min gestimuleer aan die rus; aangetrek tot intensiteit, nuwigheid en hoë-inzet situasies selfs voordat enige substansiegebruik plaasvind.	Die kras kom dikwels voor as diepgaande vlakheid en verveling eerder as angs – 'n soort grijs, onderstimuleerde leegte. Motiveringsverlies is dikwels die mees prominente klagte. Verlangens fokus dikwels op die begeerte om weer iets intens te voel, nie spesifiek op die ontsnapping van nood nie.
Met/Met ('Worrier' / 'Sage') — slow dopamine clearance, hoër basisdopamine en streschemikaltoon	Often runs anxious or overthinking at rest even before any substance use; the nervous system tends to run 'hot'.	Die kras neig om te presenteer met meer angs, rumineer en agitering wat op die vlakheid gelê word – nie net lae gemoedstoestand maar 'n bedraad-en-moeg gevoel nie. Hierdie profiel is ook meer geneig tot die hypervigilansie beskryf in Afdeling 4, en mag die slaaptekorteffekte in Afdeling 5 harder raak weens 'n reeds verhoogde stres-chemiese basislyn.
Heterozygous (Val/Met)	An intermediate baseline between the two profiles above.	Often a mix of both patterns — some flatness, some anxiety — without either being as extreme as the homozygous profiles.

GABRA2 — Die ‘Remme’ Stelsel en Hoekom Stop Met Nie Vir Almal Eenvoudig Is

Methamphetamine self is nie 'n GABA-A dwelm soos benzodiazepines — maar GABRA2 status maak steeds saak hier, vir 'n ander rede. GABRA2 regeer die digtheid en sensitiewe van die brein se primêre inhiberende ('rem') stelsel — die algemene vermoë om die remme op impuls, drang en gedrag toe te pas sodra dit reeds in beweging is. Dit is direk relevant vir hoe hard 'net stop' eintlik is vir 'n gegewe persoon, onafhangend van wilskrag.

GABRA2 Profile	What It Means for Stopping
Lower-risk / typical GABRA2 function	More baseline inhibitory 'brake' capacity available — the impulse to use, once triggered, is somewhat easier to interrupt before it becomes an action.
High-risk GABRA2 variant (thinner basaaline inhibitory tone)	Minder beknopingskap beskikbaar voor gebruik ooit begin – wat beteken die gaping tussen begeerte en aksie is werklik korter en moeiliker om slegs deur wilskrag te verbreed. Dit is 'n hardewareverskil, nie 'n karakterverskil nie, en dit is dieselfde GABRA2-argitektuur gedokumenteer in die Recovery Kernel se GABA Braking Systems materiaal vir benzodiazepines en alkohol – dit pas hier ook toe, net op algemene impulsbeheer eerder as op 'n spesifieke GABA-A-middel.

SANCTUARY

As jy voel dat dit vir jou moeiliker was om te stop as wat dit vir iemand anders voel wat weet wie dieselfde dwelm op dieselfde hoeveelheid gebruik het – daardie verskil mag werklike hardeware wees, nie 'n gat in poging of toewyding nie. GABRA2 status is presies die soort ding wat 'net stop' 'n wild verskillende taak maak vir twee verskillende senuunstelsels wat met dieselfde hunkering te doen het.

Beyond the Genes You're Born With: Does Meth Use Change Gene Expression?

Everything in the table above concerns the genetic hardware someone is born with. A separate and increasingly well-studied question is whether methamphetamine use itself changes how genes are expressed — not the underlying DNA sequence, which use does not alter, but the epigenetic marks that determine which genes get switched on or off, and how strongly. This distinction matters: nobody's genetic code is rewritten by using a substance, but the activity level of specific genes — particularly the dopamine-pathway and stress-response genes already discussed throughout this section — can genuinely shift with sustained use.

What's well-established in addiction neuroscience broadly (this is not meth-specific research alone, but methamphetamine is included in the broader literature it's drawn from): repeated stimulant exposure drives the accumulation of a transcription factor called ΔFosB in the brain's reward circuitry, which in turn alters the expression of a range of downstream genes involved in reward sensitivity and craving — this is one of the most replicated findings in addiction molecular biology, described in the foundational literature this compendium's own bibliography cites (Nestler, 2001, on the molecular basis of long-term plasticity underlying addiction). Whether methamphetamine produces additional, drug-specific epigenetic signatures beyond this general mechanism — and whether any of these changes could be passed on transgenerationally, as some animal research on other stressors and substances has suggested — is a genuinely active and not yet fully settled area of research. It would be overstating the science to present

that second part as confirmed; it's an open question worth watching, not a documented fact yet.

🛡 SANCTUARY

What's worth holding onto here: this same epigenetic mechanism runs both ways. The Recovery Kernel's own material on exercise-driven neuroplasticity describes 'the epigenetic feedback loop' directly—deliberate, matched physical activity produces measurable changes in gene expression on a documented timeline (habit and identity shift by months 1-3, measurable epigenetic changes by months 3-6, a new neurological baseline by month 6 and beyond). Whatever gene-expression changes are produced, the Tier 3 recovery work in Section 8 is not just symptom management—it's working through the identical biological channel, in the opposite direction.

Wie hierdie gids nie veilig kan maak nie: Die Geneties Kontra-geïndikeerde Profiel

Almost everything in this guide is written on the assumption that recovery, done well, is available to anyone regardless of genetic profile — the timeline may be harder or easier depending on the hardware, but it's available. There is one profile where that assumption doesn't hold, and it's worth naming plainly rather than leaving it implied inside a genetics table.

🚨 GROOT RISIKO

Vir een spesifieke genetiese en psigiatriese kombinasie dra metamfetamienegebruik 'n risiko met zich mee wat van aard verskil, nie net graad nie, van alles anders in hierdie gids – insluitend 'n werklike risiko om 'n psigotiese stoornis te ontlok wat nie ten volle oplos met abstinensie nie. Dit word hieronder volledig beskryf.

Profile	Hoekom Hierdie Een Anders Is In Soort, Nie Net Graad Nie
COMT Met/Met ('Sage'/'Worrier' — slow dopamine clearance, high baseline dopamine and stress-chemical tone) gekombineer met 'n persoonlike of gesinsgeskiedenis van psigose, skizofrenie, of bipolêre stoornis	This is the same logic the Recovery Kernel's own material applies to the most potent synthetic cathinones (MDPV, alpha-PVP): a pure dopamine-flooding stimulant, delivered into a system that already runs high baseline dopamine tone and lacks serotonin's tempering influence, produces the highest documented psychosis-precipitation risk in the stimulant category. For most people, stimulant-induced psychosis — while serious — resolves with abstinence over days to weeks. For this specific combination, the risk is meaningfully different: precipitating a first psychotic episode in someone with an underlying vulnerability can trigger a condition that does not fully resolve when the drug clears, because the The drug might have been the trigger for an underlying disorder rather than the sole cause of a temporary state.

Hoe om dit te identifiseer voordat dit te laat is

Full genetic testing (COMT rs4680, DRD2/ANKK1 Taq1A) is die mees presiese manier om te weet, maar dit is nie vir almal toeganklik nie, en die meer dringende en beskikbare skermingstool is 'n bloot historiese navraag — hierdie is die moeite werd om eerlik deur te gaan, ideaal voor enige gebruik, of so vroeg as moontlik in die herkenning van 'n gebruikerspatroon:

- Is there a family history of schizophrenia, bipolar disorder, or a psychotic episode of any kind — in a parent, sibling, or grandparent?
- Have u personally ever had 'n experience, on any substance including booze or cannabis, where u didn't just feel intensely different but lost yer grip on what was actually real — not just paranoid or antsy, but genuinely unsure what was going down around u?
- Do you tend to feel swamped, anxious, or overstimulated by normal amounts of stimulation even without any substances—the Met/Met 'Worrier' pattern described in Section 6 above?
- Het 'n psigiatries of dokter ooit die moontlikheid van 'n stemming- of psigotiese spektrumstoornis genoem, selfs een wat jy nie verder gevolg het of waarmee jy nie ingestem het nie?

⚠ WAARSKEUWING

Any combination of a 'yes' on the family or personal history questions with the Met/Met stimulant-sensitivity pattern is worth treating as a genuine contraindication, not a risk factor to weigh against the appeal of the drug's effects. This is the one place in this entire guide where the message isn't 'here's how to reduce the danger' — it's 'this specific combination is why the danger can't be fully reduced.' If this describes you or someone you're trying to help, a psychiatric evaluation before any further use — not after a crisis — is the responsible next step, and it's a 'Ek wil verstaan wat my persoonlike psigose-risiko is voordat ek dit weer gebruik.'

Phenotyping Without Test Results

- Het stimuleringsmiddels laat jou voel normaal, gefokus en lewend op 'n manier wat niks anders ooit gedoen het nie? → Waarskynlik Warrior COMT profiel met 'n DRD2 tekort daaronder.
- Het jy stimulerende middels gebruik spesifiek om te presteer – harder werk, langer, meer bereik? → Waarskynlik Val/Val COMT.
- Did u start with a prescription stimulant and escalate from there? → Probably a DRD2 deficit with tolerance developing.
- Struggle ook met depressie of emosionele vlakheid selfs wanneer nie besig om te gebruik nie? → Waarskynlik MTHFR betrokke — metileringondersteuning (5-MTHF) is 'n prioriteit saam met alles anders in Afdeling 8.

Section 7 — Use Pattern Classification

Pattern	Typical Profile	Withdrawal Severity	Recommended Posture
Laag	Occasional, non-binge use; by mouth or occasional snorting; used for a few weeks in total	Lower — a crash is real but usually brief and less bad	Stopping it is safe to do on your own, but a support contact is still worth it given the mood dip. Section 0's safety signs still apply.
Medium	Regular use, not in extended binges; weeks to a few months of use	Moderate — expect die volle kras en put fases in Afdeling 2 se tydlyn	'n Ondersteuningsstruktuur en die Seksyen 8 aanvullingsstapel vanaf dag een verminder die ernstigheid betekenisvol.
Hoog	Binge-crash cycling — multi-day use periods followed by extended crashes — for months	Hoog – die binge-crash patroon self versnel dopamienstelseluitputting verder as wat gereelde gebruik produseer.	Mediese ondersteuning tydens die krasfase word sterk aanbeveel, veral vir die suïsidie-risiko venster in Dae 1–5. Moenie dit alleen deurgaan nie.
Kroniek / IV of gerookte roete, langterm	Maande tot jare van gebruik, veral deur rook of injectie (vinniger, intenser dopamien-spikes en ooreenstemminglik dieper uitputting)	Highest — this is the population the 12–17 month D2 Recovery Kernel timeline in Section 3 was measured in	Full medical and psychiatric support during the acute crash is strongly recommended. This is also the population most likely to benefit from formal treatment engagement rather than a purely self-directed protocol.

Section 8 — Die Herstelprotokol

⚠ WAARSKEUWING

Nothing here treats suicidal ideation, psychosis, or a cardiac event. If any Section 0 sign is present, that takes priority over this protocol.

Tier 1 — Biologiese Ondersteuning

Two additions below speak directly to the glutamate side of this recovery, alongside NAC — meth's dopamine surge co-releases glutamate in cortico-striatal circuits, and excess NMDA receptor activation from that glutamate is a real contributor to the neurotoxicity in Section 3, and plausibly to the intensified, sometimes disturbing dreams that show up as REM sleep rebounds once real sleep returns (Section 4).

Support	Role
Tyrosine (500-1000mg, in the morning, away from protein)	Dopamine precursor — start once eating regularly, usually Day 3-5.
NAC (600mg twee keer per dag met kos)	Dual-action: glutamate homeostasis and craving reduction via the cystine-glutamate exchange system, AND a glutathione precursor supporting liver detoxification — chronic meth use, especially combined with any adulterants in the supply, places real burden on hepatic detox pathways, and NAC replenishes the liver's primary antioxidant molecule.
NAD+ (nicotinamide adenine dinucleotide)	Chronic substance use depletes NAD+ significantly, a coenzyme central to cellular energy production and DNA repair—part of why early recovery involves such profound fatigue. IV infusion is the most effective and most studied delivery route for acute withdrawal support specifically, with some clinical programmes reporting meaningful reduction in withdrawal discomfort; nasal spray, sublingual, and oral precursor (NMN/NR) forms are more accessible but less direct. This is a genuine option, not a first-line necessity—the foundational work below (sleep, exercise, nutrition) accounts for the large majority of physiological recovery; NAD+ and other advanced interventions are a real but a smaller addition on top of that foundation, not a substitute for it.
Calcium D-glucarate (200-400mg dae)	A complement to NAC's role above, working on a different part of liver detoxification: it inhibits beta-glucuronidase, an enzyme that can reverse the liver's Phase II conjugation process and let already-packaged toxins and hormone metabolites get reabsorbed back into circulation instead of properly excreted. Where NAC supports the glutathione conjugation pathway directly, calcium D-glucarate helps make sure what the liver has already processed actually leaves rather than recirculating — relevant background support during a period when the liver is managing both the substance itself and whatever adulterants were in the supply.

Support	Role
Magnesium glycinate (300-400mg elementeel, aand)	Magnesium sits right inside the NMDA receptor channel and physically blocks excess calcium influx unless the cell is sufficiently depolarised — it is the body's own endogenous regulator of exactly the glutamate/NMDA overactivation described above. This is on top of, not instead of, its role in Section 8's sleep support below.
P5P (pyridoxal-5-phosphate — die aktiewe vorm van witamin B6, 20-50mg daagliks)	P5P is die vereiste kofaktor vir glutamaatdekarboksilase (GAD), die ensiem wat glutamaat ombeeld in GABA. Sonder voldoende P5P is daardie omkering minder doeltreffend, wat relatief meer glutamaat en minder van die inhiberende GABA-teëwaksel oorbly – direk relevant vir die glutamaat-ryke skildery wat hierdie afdeling beskryf. Gebruik die aktiewe P5P-vorm eerder as standaard B6 (pyridoxine) indien metilering 'n kommer is, volgens die MTHFR-ry tevore.
Die 'Brain Stack' — Uridien + DHA + CDP-Choline	Direkty target op die D2-reseptor-digtheid-opregulering – die strukturele herstel van die dopamienstelsel self, nie net simptoomverligting nie.
5-MTHF (methylated folate)	Non-negotiable foundational methylation support, especially crucial if the MTHFR-linked depression/flatness pattern in Section 6 applies.
Elektrolyte en hidrasie, proteïenryke kos sodra die eetlus terugkom	Direkty ondersteun die krasfase in Afdeling 2; die liggaam se voedingsdepletasie na 'n binge is reëel en moet net so ernstig geneem word soos die psigologiese simptome.

⚠ WAARSKEUWING

On ketamine specifically, since it comes up naturally alongside glutamate: there is genuine research interest in ketamine for stimulant use disorder, working through NMDA-related synaptic plasticity rather than by 'reversing' toxicity directly, similar to its mechanism in treatment-resistant depression. It is Emerging-tier evidence, specialist-supervised only, and worth real caution rather than pursuit here specifically — the same Reward Deficiency Syndrome vulnerability discussed throughout Section 6 that made meth so compelling can make a second dissociative substance compelling too. Route of administration matters a real amount here: IV ketamine and esketamine (Spravato, intranasal, FDA-approved for treatment-resistant depression — an off-label use if applied to stimulant use disorder specifically) have far higher and more reliable bioavailability than oral ketamine, including troches — oral routes lose significant drug to first-pass liver metabolism, producing a less predictable effect. In this early, still-actively-repairing window, the more reliable IV or intranasal routes are generally considered the more clinically worthwhile option if this path is pursued at all; oral ketamine is more commonly discussed as a later-stage option, closer to the one-year mark once the acute dopamine and glutamate repair work in Section 3 is substantially further along. This is a conversation for a psychiatrist familiar with Both stimulant use disorder and ketamine's research base, not a self-directed addition to this stack.

Tier 2 — Mediese en Psigiatrië Ondersteuning

- There's no FDA-approved medication specifically for meth withdrawal — unlike the benzo or SSRI guides, this tier is about monitoring and support rather than a pharmacological bridge.
- 'n psigiatriesevaluasie tydens die krasfase is gepas indien depressie ernstig is, gegewe die werklike suïsidie-risikovenster in Dae 1-5 — dit is nie 'n oorereaksie nie, dit is 'n proporsionele reaksie op 'n gedokumenteerde risikoperiode.
- As soon as a prescriber introduces any medication (occasionally bupropion is used off-label for stimulant use disorder), disclose CYP2D6/CYP2B6 status if known, per Section 6.

Tier 2.5 — 'Ghost Scaffolding': Psychiatrist-Supervised Stimulant Stabilization

This deserves its own honest treatment rather than a single bullet, because it's a real approach that exists in specialist addiction medicine, and it's also genuinely easy to misunderstand as 'just get a prescription for the same kind of drug.' It is not that, when done correctly — but the distinction matters enough to walk through carefully.

⚠ WAARSKEUWING

This is not 'n self-directed strategy. It requires a psychiatrist or addiction medicine specialist who is actively managing the decision, monitoring for misuse risk, and treating it as a time-limited clinical tool — not an open-ended replacement. Pursuing a stimulant prescription informally, or from a general prescriber unaware of the full history, is a materially different and much riskier thing than what's described below.

The clinical logic: die Grijs en Stadige Terugkeer fases in Afdeling 2 skep werklike, meetbare uitvoerende funksiebeperking — aandag, werkgeheue, taakinisiasie — wat baie op ADHD kan lyk, of dit nou voor meth-gebruik begin het of nie. In sommige spesialistsettings (dit bestaan as gedokumenteerde praktyk in sommige lande se verslawingsmedisyneprogramme, soms genoem psiestimulanssubstitusie of agonis-styl stabilisering), word 'n deeglik bewaakte, langwerkende voorskrifstimulans — verlengde-vrystelling amfetamien-soute (Adderall XR) of soortgelyk — korttermyn gebruik spesifiek om daardie uitvoerende funksie-lek te oorbrug terwyl die dopamienstelsel in Afdeling 3 sy stadiger onderliggende herstelwerk doen, in die the same structural spirit as using buprenorphine to stabilise an opioid-dependent system rather than letting it recover entirely unsupported.

What This Is	What This Isn't
A time-limited, specialist-supervised clinical tool aimed at the specific executive-function gap in Section 2's	A green light to self-medicate the crash with any stimulant, prescribed or otherwise

What This Is	What This Isn't
Grey/Slow Return phases	
Used alongside, not instead of, the Tier 1 and Tier 3 work in this section	A substitute for the exercise, sleep, and neuroplasticity work that actually drives the DRD2 receptor recovery in Section 3
Evidence-limited and genuinely debated within addiction medicine — evidence tier: Emerging, not Established	'n Gestigde, eerste-lyn standaard van sorg — die meeste verslawingsspesialiste nog gryp hierna versigtig vir, indien al.

BEING HEARD

As die mist en vlakheid van vroeë herstel jou laat wonder of jy eintlik ADHD het, of altyd gehad het — dit is 'n legitieme vraag wat met 'n psigiatriese dokter opgewek moet word, nie iets om af te haal as net 'terugval praat' nie. Soms is dit terugval. Soms was 'n regte, voorheen ongediagnoseerde ADHD deel van wat stimulanusse so boeiend gemaak het in die eerste plek, volgens Seksie 6 se fenotipering vrae. 'n Spesialis kan jou eintlik help om die verskil te bepaal, wat presies die gesprek is wat 'ghost scaffolding' bedoel om te open, nie te vervang nie.

Tier 3 — Nervous System and Neuroplasticity Support (Die Herstel Seignaal)

- Oefening – die belangrikste ingreep in hierdie hele protokol. BDNF, die herstelsein vir die beskadigde dopamienaksone wat in Afdeling 2 en Afdeling 3 gedokumenteer word, word primêr gegenereer deur fisiese aktiwiteit. Begin so gou moontlik, selfs kortliks, en bou op na 30-45 minute van matige tot hoë intensiteit die meeste dae.
- Slaap – 7–9 ure, konsekwente rooster; BDNF sintese en geheugenkonsolidasie piek albei tydens slaap, wat beteken dat onkonsekwente slaap direk die herstelproses wat in Afdeling 3 beskryf word, vertraag.
- Morning sunlight — 20+ minutes outside, ideally within the first two hours of waking, for circadian alignment and mood support.
- A novel, challenging learning activity — 30+ minutes daily of something actively difficult (an instrument, a language, a complex skill) drives neuroplasticity more than passive activities.
- Structured daily routine and environmental redesign — removing or altering triggers tied to use locations, people, and paraphernalia.

Section 9 — On Stigma

Methamphetamine dra word 'n spesifieke, swaarder las van sosiale stigma as die meeste ander stowwe wat in hierdie biblioteek gedek word — die 'tweaker' stereotype, aannames oor kriminaliteit of morele mislukking, en 'n vlak van afkeer hoe van die drug en sy gebruikers publiek gepraat word wat alkohol- of selfs

opioïedependensie dikwels nie op dieselfde graad dra nie. Dit is dit werd om direk te noem eerder as om dit ongesproke te laat, want dit het 'n regte, gedokumenteerde praktiese effek: mense met methamphetamine-gebruikstoornis rapporteer — en navorsing oor gesondheidsorgverskaffervooroordeling bevestig — dat hulle meetbaar swakker kwaliteit sorg ontvang, minder deeglike pynbestuur, en meer afwysende behandeling. than from medical providers than people with other substance use disorders, even when presenting with the same symptoms.

BEING HEARD

As jy gevoel het of jy moes jou voorberei voordat jy vir 'n dokter, tandarts, of die ER vertel wat eintlik gebeur het—daardie instink is nie paranoia nie. Dit is 'n redelike reaksie op 'n ware patroon. Dit beteken nie dat elke verskaareder jou so sal behandel nie, en dit beteken ook nie dat jy die inligting moet verberg wat jou veilig hou (volgens Afdeling 0 en Afdeling 5 se psigose-risiko oorvlewing, 'n akkurate geskiedenis is belangrik vir jou eie sorg) — maar die versigtigheid wat jy na daardie gesprek bring, is nie 'n ooreksam op niks nie.

- As soon as a provider's tone shifts noticeably once methamphetamine use is disclosed, that's worth naming directly to them ('I want to make sure this history doesn't change the quality of care I'm getting here') or, if that doesn't work, worth escalating to a different provider — you're allowed to do that.
- Addiction medicine specialists and harm-reduction-oriented clinics generally have less of this bias baked in than general practice or emergency settings — worth seeking out specifically for ongoing care where access allows.
- Die stigma gaan oor die dwelm se reputasie, nie oor jou spesifiek nie. Die dopamienstelsel herstel wat beskryf word in Afdeling 3 gee nie om wat enigiemand oor jou karakter vermoed nie terwyl dit daardie werk doen.

Section 10 — Emergency Recognition, Restated

⚠ WAARSKEUWING

Bel findahelpline.com of 112 onmiddellik vir: suïsidale gedagtes of 'n spesifieke plan, borspyn of irreguliere hartklop, skielike ernstige hoofpyn of beroerte tekens (gesigsvalling, armswakheid, slordige spraak), psigose (paranoid wanpersepties, halusinasies), of extreme agitering met verlies aan gedragskontrole. Vir die kras simptome alleen — uitputting, hypersomnie, depressie, hunkeringe, kognitiewe mistigheid — is hierdie verwag volgens Afdeling 2, maar 'n ondersteuningskontak en, indien ernstig, 'n dieselfde-dag mediese of psigiatriese evaluasie is steeds gepas.

Section 11 — Full Protocol Cross-Reference Map

Reference	What It Offers Here
Recovery Kernel Meth/Stimulant Guide, Section 3 (The Detox Protocol)	Die volledige bronlyn wat hierdie gids se Afdeling 2 gebou word, pluss die volledige aanvullingsdosering en tydsberekening agter Afdeling 8 se Tier 1.
Recovery Kernel Compendium, Section 16.4 (Receptor Recovery — The Evidence)	The Volkow et al. and Heinz et al. citations behind Section 3's 12-17 month timeline, plus the comparative data for alcohol and cocaine receptor recovery.
Recovery Kernel Compendium, Protocol 01 (Shelter)	Die krisisstabiliseringsraamwerk wat deur die hele Afdeling 2 se kruip- en putfases verwys word.
Recovery Kernel Compendium, Protocol 05 (Environmental Redesign)	Die aanleidingverwydering en omgewingsmodifikasie-werk verwys na in Afdeling 8, Vlak 3 – identifisering en verandering van gebruikgeassosieerde mense, plekke en voorwerpe.
Recovery Kernel Compendium, Protocol 06 (Daily Structure)	Die konsekwente roetine raamwerk wat gepas begin word rond die Weeke 2-4 merk in Afdeling 2, sodra die akute kras verby is.
Recovery Kernel Compendium, Protocol 20 (Identity Reconstruction) and C-28 ('The Recast')	Gee dit aan begin rond Maande 1-3 in Afdeling 2 – nie vroeër nie. Bou 'n identiteit en 'n stel ware, hoë-intensiteit uitlaatkleppe wat geskik is vir die Warrior COMT profiel in Afdeling 6, eerder as een georganiseer rondom óf die dwelm of sy afwesigheid.
Recovery Kernel Compendium, Section 16 (Neuroplasticity Architecture)	Die volledige 90-dae herbou-raamwerk verwys na in Afdeling 3 – oefening, slaap, leer en sonlig protokolle in volledige detail, insluitend die weeklikse struktuur agter Afdeling 8 se Tier 3.
Recovery Kernel Chemsex Guide	As methamphetamine use occurred in a chemsex context specifically — the combined GHB/meth/mephedrone recovery timeline and unified supplement stack address the multi-substance picture this guide covers only for meth alone.
Recovery Kernel Komplekse Trauma en die Middelhouerverbinding	For this guide's own Section 3 (Substance-Trauma Co-Axis): chronic hypervigilance channeled into forced achievement is a documented trauma-driven pathway into stimulant use specifically — worth reading alongside this guide if that pattern applies.

Section 12 — Closing

SANCTUARY

Die dopamienstelsel wat methamphetamine oorneem het, is nie permanent gebreek nie. Die data is spesifiek en reg: meetbare herstel binne weke, substansiële strukturele herstel binne 'n jaar tot 'n jaar en 'n half, en 'n 90-dae venster waar doelbewuste oefening, slaap en nuwe uitdaging meer herbouwerk doen as enige ander punt in die proses. Die ellende van die kras was nooit 'n teken dat iets verkeerd gegaan het nie. Dit was, en is, die teken dat herstel reeds begin het.

END OF GUIDE — DISCONTINUANCE PROTOCOL: METHAMPHETAMINE

Harm Reduction Resource — Not Medical Advice — Not a Substitute for Emergency Care

Crisis Resources: 988 (Suicide & Crisis Lifeline, call or text) | Crisis Text: HOME to 741741 | SAMHSA: 1-800-662-4357